



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

THE INTRINSIC EQUATION OF A CURVE IN POLAR CO-ORDINATES.

By G. B. M. ZERR, A. M., Professor of Sciences, Texarkana, Texas.

The intrinsic equation of a curve has never been treated in polar co-ordinates in works on the subject. It is the aim of this paper to treat this subject briefly and to show that this method is somewhat simpler than the method of rectangular co-ordinates.

Let S denote the length of an arc of the curve $r=f(\theta)$ measured from some fixed point, φ the inclination of the tangent at the variable extremity to the tangent at the fixed point.

Let $y=f(x)$, then $\frac{dy}{dx}=f'(x)=-\cot \varphi$ $x=r \cos \theta$, $y=r \sin \theta$

$$\frac{dx}{d\theta} = \cos \theta \frac{dr}{d\theta} - r \sin \theta, \quad \frac{dy}{d\theta} = \sin \theta \frac{dr}{d\theta} + r \cos \theta.$$

$$\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{\sin \theta \frac{dr}{d\theta} + r \cos \theta}{\cos \theta \frac{dr}{d\theta} - r \sin \theta} = -\cot \varphi.$$

$$\therefore r \sin \theta \cot \varphi - \cos \theta \cot \varphi \frac{dr}{d\theta} = \sin \theta \frac{dr}{d\theta} + r \cos \theta.$$

$$\therefore \frac{dr}{d\varphi} = r \tan (\theta - \varphi) = f'(\theta)$$

$$\therefore f'(\theta) = f(\theta) \tan (\theta - \varphi) \dots A.$$

From A , θ is known in terms of φ , and $r = f(\varphi)$.

$$\text{Then } \frac{d\theta}{d\varphi} = F'(\varphi).$$

$$\text{Now } \frac{ds}{d\theta} = \left\{ r^2 + \left(\frac{dr}{d\theta} \right)^2 \right\}^{\frac{1}{2}} = \left[\left\{ f(\theta) \right\}^2 + \left\{ f'(\theta) \right\}^2 \right]^{\frac{1}{2}}$$

$$\therefore \frac{ds}{d\theta} = \left\{ r^2 + r^2 \tan^2 (\theta - \varphi) \right\}^{\frac{1}{2}} = r \sec (\theta - \varphi) = f(\theta) \sec (\theta - \varphi).$$

$$\frac{ds}{d\varphi} = \frac{ds}{d\theta} \cdot \frac{d\theta}{d\varphi} = F'(\varphi) f(\theta) \sec (\theta - \varphi) = r F'(\varphi) \sec (\theta - \varphi) \dots (B).$$

From B by substituting the value of r in terms of θ , and the value of θ , as found in A , in terms of φ , S may be found in terms of φ by integration.

I. Required the intrinsic equation of the circle.

The polar equation is $r=a$.

$$\therefore \frac{dr}{d\theta} = 0 = r \tan (\theta - \varphi). \quad \therefore \theta - \varphi = 0, \text{ or } \theta = \varphi \dots (1).$$

$$\frac{d\theta}{d\varphi} = 1, \therefore \frac{ds}{d\varphi} = a \sec (\varphi - \varphi) = a \sec 0 = a. \quad \therefore S = a\varphi.$$

Or we may have proceeded thus:

$$\frac{ds}{d\theta} = \left\{ r^2 + \left(\frac{dr}{d\theta} \right)^2 \right\}^{\frac{1}{2}} = a, \quad \therefore S = a\theta, \text{ but } \theta = \varphi, \quad \therefore S = a\varphi.$$

Taking the equation $r = 2a \cos \theta$ $\frac{dr}{d\theta} = -2a \sin \theta = r \tan (\theta - \varphi)$.

$$\therefore -2a \sin \theta = 2a \cos \theta \tan (\theta - \varphi).$$

$$\therefore -\tan \theta = \tan (\theta - \varphi), \therefore \theta = \frac{\varphi}{2} \dots (2).$$

$$\frac{d\theta}{d\varphi} = \frac{1}{2}, \quad \frac{ds}{d\varphi} = \frac{1}{2} \cdot 2a \cos \theta \sec (\theta - \varphi) = a \cos \frac{\varphi}{2} \sec \frac{\varphi}{2} = a,$$

$$\frac{ds}{d\varphi} = a, \quad S = a\varphi, \quad \text{or } \frac{ds}{d\theta} = \left\{ r^2 + \left(\frac{dr}{d\theta} \right)^2 \right\}^{\frac{1}{2}} = 2a, \quad S = 2a\theta, \text{ but } \theta = \frac{\varphi}{2}, \therefore S = a\varphi.$$

II. Required the intrinsic equation of the parabola. The polar equation is $r = \frac{2a}{1 + \cos \theta} = \frac{a}{\cos^2 \frac{\theta}{2}}$.

$$\frac{dr}{d\theta} = \frac{2a \sin \theta}{(1 + \cos \theta)^2} = \frac{a \sin \frac{\theta}{2}}{\cos^3 \frac{\theta}{2}} = r \tan (\theta - \varphi) = \frac{a \tan (\theta - \varphi)}{\cos^2 \frac{\theta}{2}}.$$

$$\therefore \tan \frac{\theta}{2} = \tan (\theta - \varphi), \quad \therefore \theta = 2\varphi \dots (3).$$

$$\frac{d\theta}{d\varphi} = 2, \quad \frac{ds}{d\varphi} = \frac{2a}{\cos^2 \frac{\theta}{2}} \sec (\theta - \varphi) = \frac{2a}{\cos^3 \varphi}.$$

$$\therefore S = \frac{a}{2} \log \left(\frac{1 + \sin \varphi}{1 - \sin \varphi} \right) + \frac{a \sin \varphi}{1 - \sin^2 \varphi} \quad \text{or thus}$$

$$\frac{ds}{d\theta} = \frac{a}{\cos^3 \frac{\theta}{2}}, \quad \therefore S = \frac{a}{2} \log \left(\frac{1 + \sin \frac{\theta}{2}}{1 - \sin \frac{\theta}{2}} \right) + \frac{a \sin \frac{\theta}{2}}{1 - \sin^2 \frac{\theta}{2}}, \quad \text{but } \theta = 2\varphi.$$

$$\therefore S = \frac{a}{2} \log \left(\frac{1 + \sin \varphi}{1 - \sin \varphi} \right) + \frac{a \sin \varphi}{1 - \sin^2 \varphi}.$$

III. Find the intrinsic equation to the Cardioid. The polar equation is

$$r = a(1 + \cos \theta) = 2a \cos^2 \frac{\theta}{2}, \quad \frac{dr}{d\theta} = -2a \sin \frac{\theta}{2} \cos \frac{\theta}{2} \\ = r \tan (\theta - \varphi) = 2a \cos^2 \frac{\theta}{2} \tan (\theta - \varphi).$$

$$\therefore -\tan \frac{\theta}{2} = \tan (\theta - \varphi), \quad \therefore 2\varphi = 3\theta, \quad \frac{ds}{d\theta} = \left\{ r^2 + \left(\frac{dr}{d\theta} \right)^2 \right\}^{\frac{1}{2}}$$

$$= 2a \cos \frac{\theta}{2}, \quad \therefore S = 4a \sin \frac{\theta}{2} = 4a \sin \frac{\varphi}{3}.$$

IV. Find the intrinsic equation to the curve

$$r^{\frac{1}{m}} = a^{\frac{1}{m}} \cos \frac{\theta}{m} \quad \text{or} \quad r = a \left(\cos \frac{\theta}{m} \right)^m. \quad \frac{dr}{d\theta} = -a \sin \frac{\theta}{m} \left(\cos \frac{\theta}{m} \right)^{m-1}$$

$$= r \tan (\theta - \varphi) = a \left(\cos \frac{\theta}{m} \right)^m \tan (\theta - \varphi).$$

$$\therefore -\tan \frac{\theta}{m} = \tan (\theta - \varphi). \quad \therefore (m+1)\theta = m\varphi. \quad \frac{ds}{d\theta} = \left\{ r^2 + \left(\frac{dr}{d\theta} \right)^2 \right\}^{\frac{1}{2}}$$

$$= a \left\{ \left(\cos \frac{\theta}{m} \right)^{2m} + \left(\sin \frac{\theta}{m} \right)^2 \left(\cos \frac{\theta}{m} \right)^{2m-2} \right\}^{\frac{1}{2}} = a \left(\cos \frac{\theta}{m} \right)^{m-1}.$$

$$S = a \int \left(\cos \frac{\theta}{m} \right)^{m-1} d\theta = \frac{ma}{m+1} \int \left(\cos \frac{\varphi}{m+1} \right)^{m-1} d\varphi.$$

Let $n = \frac{1}{m}$, then $S = \frac{a}{n+1} \int \left(\cos \frac{n\theta}{n+1} \right)^{\frac{1}{n}-1} d\varphi$ is the intrinsic equation to the curve $r^n = a^n \cos n\theta$.

V. Find the intrinsic equation to the Logarithmic Spiral. The polar equation is $r = be^{\frac{\theta}{c}}$ or $r = ba^{\frac{\theta}{c}}$, $\log \frac{r}{b} = \frac{\theta}{c}$, $\frac{dr}{d\theta} = \frac{r}{c} = r \tan (\theta - \varphi)$.

$$\therefore \tan (\theta - \varphi) = \frac{1}{c}, \quad \theta = \tan^{-1} \frac{1}{c} + \varphi = d + \varphi.$$

$$\frac{ds}{d\theta} = \left\{ r^2 + \left(\frac{dr}{d\theta} \right)^2 \right\}^{\frac{1}{2}} = \left(\frac{r^2 d\theta^2 + dr^2}{d\theta^2} \right)^{\frac{1}{2}} = \frac{\sqrt{1+c^2} dr}{d\theta}.$$

$$\therefore ds = \sqrt{1+c^2} dr, \quad S = \sqrt{1+c^2} \cdot r = \sqrt{1+c^2} be^{\frac{\theta}{c}} = \sqrt{1+c^2} be^{\frac{d+\varphi}{c}}.$$

The intrinsic equation for the evolute and involute can be found in the usual way.

